

# Singlet Oxygen Quenching Effect of Andaliman (*Zanthoxylum acanthopodium* DC.) Extracts in Light-Induced Lipid Oxidation

Edi Suryanto<sup>1</sup>, H. Sastrohamidjojo<sup>3</sup>, Sri Raharjo<sup>2</sup>, and Tranggono<sup>2</sup>

<sup>1</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences,  
Sam Ratulangi University, Manado

<sup>2</sup>Department of Food and Agricultural Product Technology, Faculty of Agricultural Technology

<sup>3</sup>Department of Chemistry, Faculty of Mathematic and Natural Sciences, Gadjah Mada University,  
Yogyakarta

## ABSTRACT

*Addition of andaliman extract has been shown to provide increased protection from lipid oxidation during cooking. Although andaliman extract has reported as an effective antioxidant in autoxidized system, no study have been published on its effects on light-induced oxidation or photosensitized oxidation. The objective of this research is to determine singlet oxygen quenching effects of andaliman extract on erythrosine-sensitized photooxidation of linoleic acid and in palm oil. Freeze dried ground andaliman fruit was sequentially extracted first using hexane, followed by acetone, and finally the residue was extracted by ethanol. Each of these three extracts were added to light-induced lipid peroxidation in a reaction system containing linoleic acid or palm oil with the presence of erythrosine as a photo sensitizer. Upon the exposure of fluorescent light at 4000 lux singlet oxygen is presumably formed from triplet oxygen which initiate the lipid peroxidation process. The sequential extraction of andaliman resulted in ethanolic extract containing phenolic compounds consistently show antioxidative activity presumably through singlet oxygen quenching in light-induced lipid peroxidation in either linoleic acid or RBD palm oil reaction mixture containing erythrosine as a photosensitizer. The andaliman hexane and*

*acetone extracts may contain not only phenolic compounds, but also trace amount of plant pigments which resulted in stronger lipid peroxidation effect than the antioxidative effect of phenolic substances, especially at higher level of addition.*

*Keywords : Andaliman extracts, lipid peroxidation, singlet oxygen quenching*

## INTRODUCTION

The oxidation reaction of food components leads to nutritional losses, produces undesirable flavor, and other food quality deterioration which make the food not acceptable to consumers. This oxidation reaction can be initiated by either diradical triplet oxygen or non-radical singlet oxygen. The formation of singlet oxygen in food can resulted from triplet oxygen by photosensitized reactions. The formation of singlet oxygen in foods and its quenching mechanisms have been the focus of many researches in the last twenty years (Min and Boff, 2002). This is due the fact that the rate of singlet oxygen oxidation is at least a thousand times faster than the atmospheric triplet oxygen oxidation. Singlet oxygen is produced by photosensitizers in the presence of light and triplet oxygen. Chlorophylls, pheophytin, myoglobin derivatives, riboflavin, erythrosine and methylene blue are reportedly efficient photochemical sensitizers for the formation of singlet oxygen (Whang and Peng, 1988; Yang, *et*

*al.*, 2002). Rawls and Van Santen (1970) reported that singlet oxygen participated in the initiation step of oil oxidation and the reaction rate of singlet oxygen with linoleic acid is about 1450 times greater than that of triplet oxygen.

The undesirable photosensitized lipid oxidation can be reduced by quenching singlet oxygen. Tocopherols, carotenoids, and ascorbic acid can be used for the practical reduction of singlet oxygen oxidation of oils and other oil-soluble components (Lee *et al.*, 1997). Tocopherols can also act as antioxidants in autocatalytic conditions but their singlet oxygen-quenching abilities are not effective as the carotenoid. *b*-carotene is an active  $^1O_2$  quencher in soybean oil (Lee and Min, 1990). The bleached and unbleached rosemary oleoresins had a quenching effect in the soybean oil on light-sensitized oxidation (Hall and Cuppett, 1993).

Natural antioxidant compounds can be obtained from rich plant phenolics, which are generally found in spices and aromatic plants in the tropics. The use of natural products has been long used as an ancestor of generation to generation as an additive for the food industry, besides the natural product was easy to obtain, safe for consumption, and abundant. One of the potential sources of natural products from Indonesia that is not most developed, that is wild plants that are used as spices, for example andaliman fruit. This fruit has less information about its function, characteristics, and merit of a plant as a spice.

Recently, the use of spices as antioxidants in processed food is a promising alternative to the use of synthetic antioxidants. One of the potential sources of traditional spices from North Sumatra as an antioxidant is andaliman fruit (*Zanthoxylum acanthopodium* DC.). It has a strong specific odor and pungent taste like fresh citrus and a bitter taste that stimulates saliva production. As a spice, andaliman gives a flavor such as *umami*, also it is used to preserve cooked meat for a few days against rancidity and spoilage. In addition, andaliman fruit could remove the odor of fish or meat.

Wijaya (1999) found that andaliman extracts are effective in stabilizing a system of linoleic acid emulsions against oxidation. Edi Suryanto and Rorong (2001) have reported that andaliman oleoresin exhibits

antioxidant activity greater than  $\alpha$ -tocopherol in peanut oil. Addition of andaliman extract has been shown to have scavenging activity in the DPPH radical system (Edi Suryanto *et al.*, 2004). Then andaliman extract has been shown to provide increased protection from lipid oxidation during cooking. Although andaliman extract has proven to be an effective antioxidant in an autoxidized system, no study has been published on its effects on light-induced oxidation or photosensitized oxidation. In addition, the specific compounds responsible for antioxidant active components in andaliman extracts having a singlet oxygen quenching effect has not been published.

The objective of this research was to determine the singlet oxygen quenching effects of andaliman extract on chlorophyll-sensitized photooxidation of linoleic acid and in RBD palm oil.

## MATERIALS AND METHODS

### Materials

Andaliman fruit obtained from a local market at North Sumatra will be cleaned, freeze-dried, and ground to 40 mesh. Palm oil (refined, bleached, and deodorized, RBD) was obtained from Bitung, North Sulawesi. Erythrosine was purchased from a local market. Hexane, acetone, ethanol, chloroform, sodium thiosulfate, acetic acid, potassium iodide, amylum, sodium carbonate, Folin-Ciocalteu were purchased from Merck (Darmstadt, Germany).

### Sample preparation

Andaliman fruit were cleaned, freeze-dried and ground to a fine powder (100 g) were extracted sequentially with 500 mL of hexane, acetone and ethanol for 24 hours. After filtration, the residue was extracted once more with an additional 500 mL of hexane. The hexane extracts were then pooled and saved. Similarly, the acetone extract and ethanol were obtained by extracting the residue remaining after twice extraction. The solvents in the three extracts were then evaporated using a vacuum evaporator. The resulting three extracts were then weighed and stored at  $-20^\circ\text{C}$  until use.

## Determination of total phenol content of andaliman extracts

The content of total phenol in andaliman extracts was measured by Folin-Ciocalteu assay (Hung and Yen, 2002). The absorbance of extracts was read at 750 nm in a Shimadzu UV 1601 UV-Vis Spectrophotometer. Results were expressed as mg of gallic acid/100 g extract.

## Effects of andaliman extract on erythrosine-sensitized photooxidation

Procedure is method according to Lee *et al.* (1997) with minor modification. To study the effects of andaliman extract on photosensitized oxidation of linoleic acid, sample of 0, 500, 1000, and 1500 ppm (wt/vol) andaliman extract in 0,05 M of linoleic acid (for palm oil 10.0 %) were prepared in methanol that also contained 100 ppm (wt/vol) erythrosine as a photosensitizer. Samples containing 500 ppm (wt/vol)  $\alpha$ -tocopherol were used as a positive control in the system. For palm oil were prepared in a solvent mixture (acetone/methanol, 4:1, vol/vol). Ten mL of sample was transferred into a 30 mL serum bottle. The bottles were sealed airtight with Teflon septa and aluminium caps and then were placed in the light box. The light intensity at the sample level was 4,000 lux, and temperature was room temperature. The experiment was carried out in duplicate.

## Light storage conditions for the study

The method of Lee and Min (1988) was used. The light storage box consisted of two rectangular chambers : a glass chamber (60 cm x 30 cm x 50 cm) for sample storage and the wooden box (70 cm x 50 cm x 60 cm) for light sources to the glass chamber was 12 cm. Samples were placed on the wire netting which was 10 cm above the bottom of glass chamber. The light sources, four Sylvania 15 watt cool white fluorescence lamps, were placed on the 4,000 lux. The temperature of the light storage box was kept constant at room temperature.

## Determination of oxidation of oil samples

Oxidation stability of linoleic acid and palm oil was determined by measuring the peroxide value

every 5 hours according to the AOCS (1990) method. Peroxide value will be measured every time interval of 1 hours. This activity was planned to be completed in one year. Extraction of andaliman fruit and quenching effects determination will be done at Chemistry and Biochemistry Laboratory of Food and Agricultural Product Technology, Faculty of Agricultural Technology Gadjah Mada University, Yogyakarta

## Statistical analysis

Experimental data were analyzed by the analysis of variance (ANOVA) and the significant differences among means were determined by Duncan's multiple range test (DMRT) using SPSS version 10 for windows and  $p < 0.05$  was considered to be statistically significant. Analysis were performed for the response variables of peroxide value.

## RESULTS AND DISCUSSION

### Effect of solvent on yield

The extraction was done sequentially with some solvent that possesses different polarity to separate compounds in andaliman fruit. Extraction using hexane could dissolve less polar compounds, whereas acetone could dissolve semi polar compounds and the use of ethanol would recover the more polar compounds. The yield of hexane extract (HE) gave higher yields from sequential extraction than ethanol extract (EE) and acetone extract (AE). Hexane extract had the highest yield, due to the essential oil and lipid were probably extracted.

Table 1. Extraction yield and total phenol content


Andaliman extract	Yield (mg/g of andaliman extract)	Total phenol content (g/100 g of extract)
Hexane extract (HE)	78,06 $\pm$ 2,48	2,77 $\pm$ 0,58
Acetone extract (AE)	31,75 $\pm$ 5,56	9,10 $\pm$ 0,03
Ethanol extract (EE)	69,98 $\pm$ 3,36	12,53 $\pm$ 0,59

Mean value of three measurements  $\pm$  SD

Choice of solvent and extraction procedures were also considered. Presence of chlorophyll in hexane and acetone extracts was responsible for the deep-green coloration. Chlorophyll and their derivatives had been reported effective as sensitizer for production of singlet oxygen under light illumination (Rawls and Van Santen, 1970). Hexane had been considered the solvent of choice for further extraction, due to the essential oil, oleoresin and a possible presence of fatty acids of andaliman fruit can be removed by this solvent.

### Effect of erythrosine on linoleic acid

Effect of erythrosine at different concentrations on linoleic acid on peroxide value for 0.03 M of linoleic acid under light (4000 lux) exposure for 5 hours in room temperature are shown in Figure 1. Effect of concentrations at 25, 50, 75, and 100 ppm showed increased changes in peroxide value during expose to light for 5 hours ( $p < 0,05$ ). However, changes in peroxide value of without erythrosine not showed increasing significantly. This result proved that without erythrosine couldn't produced oxygen singlet of triplet oxygen, although the sample under light exposure for 5 hours.




**Figure 1.** Changes in peroxide value of linoleic acid (0,03 M) with different concentration erythrosine during exposure to light (4000 lux) for 5 hours (Er: erythrosine, K-1: light without erythrosine and K-2: dark with erythrosine).


Min and Bob (2002) had reported that singlet oxygen could be produced triplet oxygen with presence sensitizer and light. There is sensitizer as erythrosine can increase oxidation reaction, due to sensitizer possess ability for absorption of light energy. The sensitizer can transfer its energy to the most stable triplet oxygen and converting it to a higher energy level, singlet oxygen. It would then attack the double bonds in unsaturated fatty acids (Khan and Shahidi, 2002). The primary products of photooxidation is hidroperoxide, for example photooxidation on linoleic acid can produce hidroperoxide of position of conjugated double bond at 9-OOH and 13-OOH and non conjugated at 10-OOH and 12-OOH. The contrary, the autooxidation of triplet oxygen only produce conjugated hidroperoxide at 9-OOH and 13-OOH (Neff and Frangkel, 1980). This hidroperoxides are subsequently decomposed to off-flavours volatiles (Warner and Frankel, 1987; Jung *et al.*, 1991).

### Effect of andaliman extract on linoleic acid


Effect of variation concentration from hexane extract (HE), acetone extract (AE), and ethanol extract (EE) on peroxide value for 0.05 M of linoleic acid in methanol for expose to light in room temperature (Figure 2, 3, and 4). Effect of EE concentration from extracts showed significant on sample peroxide value (PV) for 5 hours exposure to light in room temperature, for example EE concentration of 500, 1000, and 1500 ppm, sample peroxide is the similar wherease at concentrations of 500, 1000, and 1500 ppm HE can enhance peroxide value. However, AE at concentration of 1500 ppm indicate increasing peroxide value. A probably explanation, photosensitizer of erythrosine may product singlet oxygen from triplet oxygen when there is exposed to light. Data of peroxide value in linoleic acid is relative similar for 5 hours to expose light without sensitizer. The formation of singhlet oxygen product by photosensitizer that accelarated lipid oxidation. Therefore, erythrosine may act as photosensitizer for accelarate of linoleic acid oxidation in model system to expose light (Yang *et al.*, 2002).



**Figure 2.** Effect of 500 ppm andaliman extract on oxidation of singlet oxygen in linoleic acid for 5 hours (VE: a-tocopherol; K-1: light with erythrosine and K-2: light without erythrosine).



**Figure 4.** Effect of 1500 ppm andaliman extract on oxidation of singlet oxygen in linoleic acid for 5 hours (VE: a-tocopherol; K-1: light with erythrosine and K-2: light without erythrosine).




**Figure 3.** Effect of 1000 ppm andaliman extract on oxidation of singlet oxygen in linoleic acid for 5 hours (VE: a-tocopherol; K-1: light with erythrosine and K-2: light without erythrosine).

Effect of concentrations present of 500, 1000, and 1500 ppm AE in model system showed that different significantly with 1000 ppm a-tocopherol. Although the differences in concentrations of 500 ppm and 1000 ppm AE is not significantly with 1000 ppm of a-tocopherol. Vitamine E (VE) is nature compound to used inhibitor in lipid oxidation for food. Therefore, a-tocopherol had reported as quencher of singlet oxygen in soy been oil (Jung *et al.*, 1991). The other concentration showed that 1500 ppm of HE can indicate prooxidative. In other hand, the changes of peroxide value in linoleic acid more increasing with different concentrations of extract. This is probably due to effect of chlorophyll catalitic that obtained to HE.


#### Effect of andaliman extract on palm oil (RBD)

The changes of eroxide value (PV) in palm oil (RBD) with different concentration of extracts are given in Figure 5, 6, and 7. Oxidation reaction of palm oil will be mesuared every time interval for 5 hours at room temperature and stored exposed to light of 4000 lux. Peroxide value increase in palm oil that addition erythrosine as photosensitizer of 45 meq/kg, whereas palm oil without sensitizer was 4.5 meq/kg.


The curve in Figure 5, 6, and 7 showed that EE at concentrations of 500, 1000, 1500 and 500 ppm of AE could be increased rate of oxidation for peroxide. However, fourth the extracts is not different with tocopherol ( $p < 0.05$ ) as the control positive in this study. At concentrations of 500, 1000, and 1500 ppm showed that increasing peroxide value and as prooxidative. This could be seen from peroxide value of HE is not different with palm oil that addition erythrosine ( $p < 0.05$ ). This is as well as for 1000 ppm and 1500 ppm of AE. This explanation caused when to extract with hexane and acetone, perhaps chlorophyll extracted in both the solvent, whereas at EE was not extracted. In this study has not done determination of chlorophyll content for every the extracts. The chlorophyll is sensitizer which further probably can act synergistically with erythrosine to generate singlet oxygen so that oxidation rate of peroxide formation is increase. Therefore, dechlorophyll of extracts as nature antioxidant resource in protect of unsaturated fatty acid could be prolong storage.



**Figure 5.** Effect of 500 ppm andaliman extract on oxidation of singlet oxygen in RBD palm oil for 5 hours (VE: a-tocopherol; K-1: light with erythrosine and K-2: light without erythrosine).



**Figure 6.** Effect of 1000 ppm andaliman extract on oxidation of singlet oxygen in RBD palm oil for 5 hours (VE: a-tocopherol; K-1: light with erythrosine and K-2: light without erythrosine).



**Figure 7.** Effect of 1500 ppm andaliman extract on oxidation of singlet oxygen in RBD palm oil for 5 hours (VE: a-tocopherol; K-1: light with erythrosine and K-2: light without erythrosine).

Effect of prooxidative from HE and AE could be caused to present chlorophyll. Endo *et al.* (1985) had reported that chlorophyll and its derivate can promote lipid oxidation for storage. Wanasundara and Shahidi (1998), in its study showed that etanol-aqueous extract possessed effect of prooxidant in edible oil, perhaps caused effect of catalitic from

chlorophyll. The other study showed that effect of photosensitization of chlorophyll present in spices. May be more important than the effect of the antioxidants for food exposed to light (Madsen *et al.* 1998 ; Wanasundara and Shahidi, 1998). This estimate that EE of obtained sequentially was not extracted chlorophyll again so that may act as quencher of oxygen singlet.

The edible oil included palm oil to content a-tocopherol and carotenoid (Gunstone, 1996). In this study does not use palm oil (RBD) that free a-tocopherol. Based on, a probably that a-tocopherol addition in palm oil of 1000 ppm to be make more effective as quencher of oxygen singlet. Addition for, it may be phenolics compound to obtain in EE and a-tocopherol are naturally in palm oil (RBD) possesse effect of synergismcally.

### CONCLUSION

The sequential extraction of andaliman fruit using hexane, acetone and ethanol resulted in three extracts with different concentrations. The ethanol extract seems to contain phenolic compound showed effect as quencher of singlet oxygen on linoleic acid and palm oil photooxidation with present erythrosine as sensitizer.

### ACKNOWLEDGEMENT

The authors would like to thank the Indonesia Toray Science Foundation for providing research grant for this study in 2003.

### REFERENCES

AOCS, 1990. *Official and Tentative Methods*. American Oil Chemists Society, Champaign, IL.

Endo, Y., R. Usuki, and T. Kaneda. (1985). Antioxidant Effects on Chlorophylls and Pheophytin on the Autooxidation of Oils in The dark II. *JAACS*. 62: 1387-1390.

Edi Suryanto, dan J.A Rorong. 2001, Isolasi Antioksidan Fenolik dari Oleoresin Buah Andaliman (*Zanthoxylum acanthopodium DC.*). *Eugenia* 7 : 88-92.

Edi Suryanto, Sri Raharjo, H. Sastrohamidjojo and Tranggono. 2004. Antiradical Activity of Andaliman (*Zanthoxylum acanthopodium DC.*) Fruit Extract. *Indonesian Food and Nutrition Progress*. 11: 16-20.

Gunstone, F.D. 1996. *Fatty Acid and Lipid Chemistry*. Blakie Academic & Professional, London.

Hall III, C and S. Cuppsett. 1993. The effects of Bleached and Unbleached Rosemary Oleoresins on Light-Sensitized Oxidation of Soybean Oil. *JAACS*. 70: 477-482.

Hung and Yen. 2002. Antioxidant Activity of Phenolic Compounds Isolated from *Mesona procumbens Hemsl.* *J. Agric Food Chem.* 50: 2993-2997.

Jung, M.Y., E. Choe, and D.B Min. 1991. Effects of a, b, g-Tocopherol on Chlorophylls Photosensitized Oxidation of Soybean Oil. *J. Food Sci.* 56: 807-815.

Khan, M.A. and F. Shahidi. Photooxidative Stability of Stripped and Non-stripped Borage and Evening Primrose Oils and Their Emulsions in Water. *Food Chemistry*. 79: 47-53.

Lee, S-H and D.B. Min. 1990. Effects, Quenching Mechanism, and Kinetic of Carotenoids in Chlorophyll-Sensitized Photooxidation of Soybean Oil. *J. Agric. Food Chem.* 38: 1630-1634.

Lee, K.H., M.Y. Jung, and S.Y. Kim. 1997. Quenching Mechanism and Kinetics of Ascorbyl Palmitate for the Reduction of the Photosensitized Oxidation of Oils. *JAACS*. 74: 1053-1057.

Madsen, H.L., B. Sorensen, L. H. Skibsted, and G Bertelsen. 1998. The antioxidative Activity of Summer Savory (*Satureja hortensis L.*) and Rosemary (*Rosmarinus officinalis L.*) in Dressing Stored Exposed to Light or in Darkness. *Food Chemistry*. 63: 173-180.

Min, D.B. and J.M. Boff. 2002. Chemistry and Reaction of Singlet Oxygen in Foods. *Comprehensive Reviews in Food Science and Food Safety*. 1: 58-72.

Neff, W.E. and E.N. Frankel, 1980. Quantitative Analysis of Hydroxystearate Isomers from Hydroperoxides by High Pressure Liquid Chromatography of Autoxidized and Photosensitized-Oxidized Fatty Esters. *Lipid*. 15: 587-590.

Rawls, H.R. and Van Santen, P.J. 1970. Possible Role of Singlet Oxidation in The initiation of Fatty Acid Autoxidation. *JAACS*. 47: 121-125.

Yang, W.T., J.H. Lee, and D.B. Min. 2002. Quenching Mechanisms and Kinetics of a-Tocopherol and b-Carotene on the Photosensitizing Effect of Synthetic Food Colorant FD&C Red No. 3. *J. Food. Sci.* 67: 507-510.

- Warner, K. and E.N Frankel. 1987. Effect of b-Carotene on Light Stability of Soybean Oil. *JAACS*. 64: 213-218.
- Wanasundara, U.N., and F. Shahidi. 1998. Antioxidant and Pro-oxidant Activity of Green Tea Extracts in Marine Oils. *Food Chemistry*. 63: 335-342.
- Whang, K. and I.C. Peng. 1988. Photosensitized Lipid Peroxidation in Ground Pork and Turkey. *J. Food Sci.* 53: 1596-1598.
- Wijaya, C.H. 1999. Andaliman Rempah Tradisional Sumut dengan Aktivitas Antioksidan dan Antimikroba. *Bul. Teknol. dan Industri Pangan*. 2: 59-61